

# Normal vs abnormal motions of the knee

To understand the functions of the knee ligaments, how they get injured, and how their injury would cause abnormal/unstable knee motions, you first need to make sure you can recognize *normal* motions of the knee. So, you'll first practice simulating the normal motions of the knee that you'll repeat later as you remove each ligament to test for abnormal motion and instability.

The sections below will walk you through simulating **anterior-posterior translation**, **longitudinal rotation**, and **varus-valgus rotation**, each at near full extension of the knee and at 90 degrees of flexion. Practice each motion multiple times with all of the ligaments in place until you get a good feel for what is the normal range of motion. Once you start removing ligaments, you can always reattach the ligament to remind yourself of the normal motion. For this entire activity, you'll hold the femur still and move the tibia relative to a static femur.

## Anterior-posterior translation

### Normal range of motion

Simulate **anterior-posterior (AP) translation** of the tibia at near full extension of the knee and at 90° of flexion, using the video below as a guide; here "near full extension" means about 20° short of full extension (about 160° between the femur and tibia).

Video showing how to simulate anterior-posterior translation at near full extension and at 90° of flexion. Note that unlike the video, your knee kit won't have muscles and/or the patella attached. That's OK—these are not needed to simulate this motion.

With all of the ligaments intact, you should only be able to translate the tibia approximately 1 cm (10 mm) in total. You can use your ruler to verify this. You can also think of this as the tibia moving 5 mm anterior and 5 mm posterior relative to its neutral/average position (i.e., when the tibia is in the middle of its motion path).

### Abnormal range of motion

You can recognize abnormal AP translation as movement of the tibia greater than 5 mm either in the anterior or posterior direction from its neutral position.


## Longitudinal rotation

### Normal range of motion

Next, simulate **longitudinal rotation** (i.e., rotation about its long-axis) of the tibia at near full extension of the knee and at 90° of flexion, using the video below as a guide.

Video showing how to simulate longitudinal rotation of the tibia at near full extension and at 90° of flexion. Note that unlike the video, your knee kit won't have muscles and/or the patella attached. That's OK—these are not needed to simulate this motion.

You don't need to worry about measuring the number of degrees the tibia can rotate in each direction. Rather, pay attention to the articular cartilage surfaces where the femur and tibia contact one another. Notice how during normal motion (and in a healthy knee) and when there is force pushing the bones at a joint together (joint under compression), there are is always articular cartilage on each side/bone at the point of contact.



When a healthy joint is under compression and during normal motion, there is always articular cartilages on opposing sides of the joint at the point of contact.

You do *not* want bone-on-cartilage or bone-on-bone contact at a joint (that's what happens with osteoarthritis—not good!). Rather, you want continuous cartilage contact for smooth (and pain-free) motion.

## Abnormal range of motion

You can recognize abnormal longitudinal rotation when the tibia rotates so far relative to the femur that there is no longer any articular cartilage at the point of contact on one or both bones.

Varus-valgus rotation

## Normal range of motion


Lastly, simulate **varus-valgus rotation** of the tibia at near full extension of the knee and at 90° of flexion, using the video below as a guide.

Video showing how to simulate varus-valgus rotation at near full extension and at 90° of flexion. Note that unlike the video, your knee kit won't have muscles and/or the patella attached. That's OK—these are not needed to simulate this motion.

As you simulate this rotation, take a close look at where the femur and tibia are contacting one another inside the joint. During varus rotation, a gap forms between the lateral condyles of the femur and tibia. During valgus rotation, a very small gap opens up between the medial condyles (hardly visible). This gap formation is called **lift off**.

Video showing lift off between the lateral femoral condyle and lateral tibial condyle during varus-valgus rotation. The knee is flexed at 90°, the view is the from the anterior, and the patella has been pulled down to show the condyles.

During varus rotation, a normal lift off between the lateral condyles is about 2 mm at near full extension and about 10 mm at 90° of flexion. During valgus rotation, the amount of lift off between the medial condyles is less than 1 mm.



An example of normal maximum lift off. In this case, between the lateral femoral condyle and lateral tibial condyle during varus-valgus rotation at 90° of knee flexion.

## Abnormal range of motion

You can recognize abnormal motion during varus or valgus rotation as a greater than normal lift off between the lateral or medial condyles, respectively.

